arXiv:1611.02595v2 [math.DG] 3 Dec 2016

AFFINE TRANSLATION SURFACES IN THE ISOTROPIC 3-SPACE

MUHITTIN EVREN AYDIN*, MAHMUT ERGUT

ABSTRACT. The isotropic 3-space $\mathbb{I}^3$ is a real affine 3-space endowed with the metric dx^2+dy^2. In this paper we describe Weingarten and linear Weingarten affine translation surfaces in $\mathbb{I}^3$. Further we classify the affine translation surfaces in $\mathbb{I}^3$ that satisfy certain equations in terms of the position vector and the Laplace operator.

1. INTRODUCTION

It is well-known that a surface is called *translation surface* in a Euclidean 3-space $\mathbb{R}^3$ if it is the graph of a function $z(x,y)=f(x)+g(y)$ for the standart coordinate system of $\mathbb{R}^3$. One of the famous minimal surfaces of $\mathbb{R}^3$ is Scherk's minimal translation surface which is the graph of the function

$$z(x,y)=\frac{1}{c}\log\left|\frac{\cos(cx)}{\cos(cy)}\right|,\ c\in\mathbb{R}^*:=\mathbb{R}-\{0\}.$$

In order for more generalizations of the translation surfaces to see in various ambient spaces we refer to [4, 5, 7, 12, 16, 19, 20, 24, 26].

In 2013, H. Liu and Y. Yu [14] defined the *affine translation surfaces* in $\mathbb{R}^3$ as the graph of the function

$$z(x,y)=f(x)+g(y+ax),\ a\in\mathbb{R}^*$$

and described the minimal affine translation surfaces which are given by

$$z(x,y)=\frac{1}{c}\log\left|\frac{\cos\left(c\sqrt{1+a^2}x\right)}{\cos(c[y+ax])}\right|,\ a,c\in\mathbb{R}^*.$$

These are called *affine Scherk surface*. Then H. Liu and S.D. Jung [15] classified the affine translation surfaces in $\mathbb{R}^3$ of arbitrary constant mean curvature.

In the isotropic 3-space $\mathbb{I}^3$, there exist three different classes of translation surfaces given by (see [18, 25])

$$\begin{cases} z(x,y)=f(x)+g(y),\\ y(x,z)=f(x)+g(z),\\ x(y,z)=\frac{1}{2}\left(f\left(\frac{y+z-\pi}{2}\right)+g\left(\frac{\pi-y+z}{2}\right)\right),\end{cases}$$

where x,y,z are the standart affine coordinates in $\mathbb{I}^3$. These surfaces are respectively called *translation surfaces of Type 1,2,3* in $\mathbb{I}^3$. Such surfaces of constant isotropic Gaussian and mean curvature were obtained in [18] as well as Weingarten ones.

2000 *Mathematics Subject Classification.* 53A35, 53A40, 53B25.

Key words and phrases. Isotropic space, affine translation surface, Weingarten surface.

The translation surfaces of Type 1 in $\mathbb{I}^3$ that satisfy the condition

$$\triangle^{I,II} r_i = \lambda_i r_i,\ \lambda_i \in \mathbb{R},\ i = 1, 2, 3,$$

were presented in [13], where r_i is the coordinate function of the position vector and $\triangle^{I,II}$ the Laplace operator with respect to the first and second fundamental forms, respectively. This condition is natural, being related to the so-called *submanifolds of finite type*, introduced by B.-Y. Chen in the late 1970's (see [8, 9, 11]). More details of translation surfaces in the isotropic spaces can be found in [2, 3, 6].

In this paper, we investigate the affine translation surfaces of Type 1 in $\mathbb{I}^3$, i.e. the graphs of the function

$$z(x, y) = f(ax + by) + g(cx + dy),\ ad - bc \neq 0$$

and classify ones of Weinagarten type. Morever we describe the affine translation surfaces of Type 1 that satisfy the condition $\triangle^{I,II} r_i = \lambda_i r_i$.

2. Preliminaries

The isotropic 3-space $\mathbb{I}^3$ is a real affine space defined from the projective 3-space $P\left(\mathbb{R}^3\right)$ with an absolute figure consisting of a plane ω and two complex-conjugate straight lines f_1, f_2 in ω (see [1, 10, 17], [21]-[23]). Denote the projective coordinates by (X_0, X_1, X_2, X_3) in $P\left(\mathbb{R}^3\right)$. Then the absolute plane ω is given by $X_0 = 0$ and the absolute lines f_1, f_2 by $X_0 = X_1 + iX_2 = 0$, $X_0 = X_1 - iX_2 = 0$. The intersection point $F(0:0:0:1)$ of these two lines is called the absolute point. The group of motions of $\mathbb{I}^3$ is a six-parameter group given in the affine coordinates $x = \frac{X_1}{X_0}$, $y = \frac{X_2}{X_0}$, $z = \frac{X_3}{X_0}$ by

$$(x, y, z) \longmapsto (x', y', z') : \begin{cases} x' = a_1 + x\cos\phi - y\sin\phi, \\ y' = a_2 + x\sin\phi + y\cos\phi, \\ z' = a_3 + a_4 x + a_5 y + z, \end{cases}$$

where $a_1, ..., a_5, \phi \in \mathbb{R}$.

The metric of $\mathbb{I}^3$ is induced by the absolute figure, i.e. $ds^2 = dx^2 + dy^2$. The lines in $z-$direction are called *isotropic lines*. The planes containing an isotropic line are called *isotropic planes*. Other planes are *non-isotropic*.

Let M be a surface immersed in $\mathbb{I}^3$. We call the surface M *admissible* if it has no isotropic tangent planes. Such a surface can get the form

$$r : D \subseteq \mathbb{R}^2 \longrightarrow \mathbb{I}^3 :\ (x, y) \longmapsto (r_1(x, y), r_2(x, y), r_3(x, y)).$$

The components E, F, G of the first fundamental form I of M can be calculated via the metric induced from $\mathbb{I}^3$.

Denote $\triangle^I$ the Laplace operator of M with respect to I. Then it is defined as

$$\triangle\phi = \frac{1}{\sqrt{|W|}}\left\{\frac{\partial}{\partial x}\left(\frac{G\phi_x - F\phi_y}{\sqrt{|W|}}\right) - \frac{\partial}{\partial y}\left(\frac{F\phi_x - E\phi_y}{\sqrt{|W|}}\right)\right\}, \tag{2.1}$$

where ϕ is a smooth function on M and $W = EG - F^2$.

The unit normal vector field of M^2 is completely isotropic, i.e. $(0, 0, 1)$. Morever, the components of the second fundamental form II are

$$L = \frac{\det(r_{xx}, r_x, r_y)}{\sqrt{EG - F^2}},\ M = \frac{\det(r_{xy}, r_x, r_y)}{\sqrt{EG - F^2}},\ N = \frac{\det(r_{yy}, r_x, r_y)}{\sqrt{EG - F^2}}, \tag{2.2}$$

where $r_{xy} = \frac{\partial^2 r}{\partial x \partial y}$, etc.

The *relative curvature* (so-called the *isotropic curvature* or *isotropic Gaussian curvature*) and the *isotropic mean curvature* are respectively defined by

$$K=\frac{LN-M^2}{EG-F^2},\ H=\frac{EN-2FM+LG}{2(EG-F^2)}. \tag{2.3}$$

Assume that nowhere M has parabolic points, i.e. $K\neq 0$. Then the Laplace operator with respect to II is given by

$$\triangle^{II}\phi=-\frac{1}{\sqrt{|w|}}\left\{\frac{\partial}{\partial x}\left(\frac{N\phi_x-M\phi_y}{\sqrt{|w|}}\right)-\frac{\partial}{\partial y}\left(\frac{M\phi_x-L\phi_y}{\sqrt{|w|}}\right)\right\} \tag{2.4}$$

for a smooth function ϕ on M and $w=LN-M^2$.

In particular, if M is a graph surface in $\mathbb{I}^3$ of a smooth function $z(x,y)$ then the metric on M induced from $\mathbb{I}^3$ is given by dx^2+dy^2. Thus its Laplacian turns to

$$\triangle^{I}=\frac{\partial^2}{\partial x^2}+\frac{\partial^2}{\partial y^2}. \tag{2.5}$$

The matrix of second fundamental form II of M corresponds to the Hessian matrix $\mathcal{H}(z)$, i.e.,

$$\begin{pmatrix} L & M\\ M & N\end{pmatrix}=\begin{pmatrix} z_{xx} & z_{xy}\\ z_{xy} & z_{yy}\end{pmatrix}.$$

Accordingly, the formulas (2.3) reduce to

$$K=\det\left(\mathcal{H}(z)\right),\ H=\frac{trace\left(\mathcal{H}(z)\right)}{2}. \tag{2.6}$$

3. Weingarten affine translation surfaces

Let M be the graph surface in $\mathbb{I}^3$ of the function $z(x,y)=f(u)+g(v)$, where

$$u=ax+by,\ v=cx+dy. \tag{3.1}$$

If $ad-bc\neq 0$, we call the surface M *affine translation surface of Type 1* in $\mathbb{I}^3$ and the pair (u,v) *affine parameter coordinates*.

In the particular case $a=d=1$ and $b=c=0$ (or $a=d=0$ and $b=c=1$), such a surface reduces to the translation surface of Type 1 in $\mathbb{I}^3$. Let us fix some notations to use remaining part:

$$\frac{\partial f}{\partial x}=a\frac{df}{du}=af',\ \frac{\partial f}{\partial y}=bf',\ \frac{\partial g}{\partial x}=c\frac{dg}{dv}=cg',\ \frac{\partial g}{\partial y}=dg',$$

and so on. By (2.6), the relative curvature K and the isotropic mean curvature H of M turn to

$$K=(ad-bc)^2 f''g'' \text{ and } 2H=\left(a^2+b^2\right)f''+\left(c^2+d^2\right)g''. \tag{3.2}$$

Now we can state the following result to describe the Weingarten affine translation surfaces of Type 1 in $\mathbb{I}^3$ that satisfy the condition

$$K_xH_y-K_yH_x=0, \tag{3.3}$$

where the subscript denotes the partial derivative.

Theorem 3.1. *Let M be a Weingarten affine translation surface of Type 1 in $\mathbb{I}^3$. Then one of the following occurs:*

(i) M is a quadric surface given by

$$z(x,y) = c_1u^2 + \frac{c_1\left(a^2+b^2\right)}{\left(c^2+d^2\right)}v^2 + c_2u + c_3v + c_4,\ c_1,...,c_4 \in \mathbb{R};$$

(ii) M is of the form either

$$z(x,y) = f(u) + c_1v^2 + c_2v + c_3,\ f''' \neq 0,\ c_1,c_2,c_3 \in \mathbb{R}$$

or

$$z(x,y) = g(v) + c_1u^2 + c_2u + c_3,\ g''' \neq 0,\ c_1,c_2,c_3 \in \mathbb{R},$$

where (u,v) is the affine parameter coordinates given by (3.1).

Remark 3.1. We point out that a *quadric surface* in $\mathbb{I}^3$ is the set of the points satisfying an equation of the second degree.

Proof. It follows from (3.2) and (3.3) that

$$\left[\left(a^2+b^2\right)f'' - \left(c^2+d^2\right)g''\right]f'''g''' = 0. \tag{3.4}$$

To solve (3.4) , we have several cases:

Case (a). $\left(a^2+b^2\right)f'' = \left(c^2+d^2\right)g''$. Then we derive

$$z(x,y) = c_1u^2 + \frac{c_1\left(a^2+b^2\right)}{\left(c^2+d^2\right)}v^2 + c_2u + c_3v + c_4,\ c_1,...,c_4 \in \mathbb{R},$$

which gives the statement (i) of the theorem.

Case (b). $\left(a^2+b^2\right)f'' \neq \left(c^2+d^2\right)g''$. Then, by (3.4), the surface has the form either

$$z(x,y) = g(v) + c_1u^2 + c_2u + c_3,\ g''' \neq 0$$

or

$$z(x,y) = f(u) + c_4v^2 + c_5v + c_6,\ f''' \neq 0,\ c_1,...,c_6 \in \mathbb{R}.$$

This implies the second statement of the theorem. Therefore the proof is completed. □

Now we intend to find the linear Weingarten affine translation surfaces of Type 1 in $\mathbb{I}^3$ that satisfy

$$\alpha K + \beta H = \gamma,\ \alpha,\beta,\gamma \in \mathbb{R},\ (\alpha,\beta,\gamma) \neq (0,0,0). \tag{3.5}$$

Without lose of generality, we may assume $\alpha \neq 0$ in (3.5) and thus it can be rewritten as

$$K + 2m_0H = n_0,\ 2m_0 = \frac{\beta}{\alpha},\ n_0 = \frac{\gamma}{\alpha}. \tag{3.6}$$

Hence the following result can be given.

Theorem 3.2. *Let M be a linear Weingarten affine translation surface of Type 1 in $\mathbb{I}^3$ that satisfies* (3.6)*. Then we have:*

(i) M is a quadric surface given by

$$z(x,y) = c_1u^2 + c_2v^2 + c_3u + c_4v + c_5,\ c_1,...,c_5 \in \mathbb{R};$$

(ii) M is of the form either

$$z(x,y) = f(u) - \frac{m_0\left(a^2+b^2\right)}{2(ad-bc)^2}v^2 + c_1 v + c_2,\ f''' \neq 0,\ c_1, c_2 \in \mathbb{R}$$

or

$$z(x,y) = g(v) - \frac{m_0\left(c^2+d^2\right)}{2(ad-bc)^2}u^2 + c_1 u + c_2,\ g''' \neq 0,\ c_1, c_2 \in \mathbb{R},$$

where (u,v) is the affine parameter coordinates given by (3.1).

Proof. Substituting (3.2) in (3.6) gives

$$(ad-bc)^2 f''g'' + m_0\left(a^2+b^2\right)f'' + m_0\left(c^2+d^2\right)g'' = n_0. \tag{3.7}$$

After taking partial derivative of (3.7) with respect to u and v, we deduce $f'''g''' = 0$. If both f''' and g''' are zero then we easily obtain the first statement of the theorem. Otherwise, we have the second statement of the theorem. This proves the theorem. □

Example 3.1. Consider the affine translation surface of Type 1 in $\mathbb{I}^3$ with

$$z(x,y) = \cos(x-y) + (x+y)^2,\ -\frac{\pi}{6} \leq x, y \leq \frac{\pi}{6}.$$

This surface plotted as in Fig. 1 satisfies the conditions to be Weingarten and linear Weingarten.

4. Affine translation surfaces satisfying $\triangle^{I,II} r_i = \lambda_i r_i$

This section is devoted to classify the affine translation surfaces of Type 1 in $\mathbb{I}^3$ that satisfy the conditions $\triangle^{I,II} r_i = \lambda_i r_i$, $\lambda_i \in \mathbb{R}$. For this, we get a local parameterization on such a surface as follows

$$\begin{aligned} r(x,y) &= (r_1(x,y), r_2(x,y), r_3(x,y)) \\ &= (x, y, f(ax+by) + g(cx+dy)). \end{aligned} \tag{4.1}$$

Thus we first give the following result.

Theorem 4.1. *Let M be an affine translation surface of Type 1 in $\mathbb{I}^3$ that satisfies $\triangle^I r_i = \lambda_i r_i$. Then it is congruent to one of the following surfaces:*

(i) $(\lambda_1, \lambda_2, \lambda_3) = (0,0,0)$

$$z(x,y) = c_1 u^2 - \frac{c_1\left(a^2+b^2\right)}{\left(c^2+d^2\right)}v^2 + c_3 u + c_4 v + c_5;$$

(ii) $(\lambda_1, \lambda_2, \lambda_3) = (0,0,\lambda > 0)$

$$z(x,y) = c_1 e^{\sqrt{\frac{\lambda}{a^2+b^2}}u} + c_2 e^{-\sqrt{\frac{\lambda}{a^2+b^2}}u} + c_3 e^{\sqrt{\frac{\lambda}{c^2+d^2}}v} + c_4 e^{-\sqrt{\frac{\lambda}{c^2+d^2}}v};$$

(iii) $(\lambda_1, \lambda_2, \lambda_3) = (0,0,\lambda < 0)$

$$\begin{aligned} z(x,y) &= c_1 \cos\left(\sqrt{\tfrac{-\lambda}{a^2+b^2}}u\right) + c_2 \sin\left(\sqrt{\tfrac{-\lambda}{a^2+b^2}}u\right) + c_3 \cos\left(\sqrt{\tfrac{-\lambda}{c^2+d^2}}v\right) \\ &\quad + c_4 \sin\left(\sqrt{\tfrac{-\lambda}{c^2+d^2}}v\right), \end{aligned}$$

where (u,v) is the affine parameter coordinates given by (3.1) and $c_1, ..., c_5 \in \mathbb{R}$.

Proof. It is easy to compute from (2.5) and (4.1) that

$$\triangle^I r_1 = \triangle^I r_2 = 0 \tag{4.2}$$

and

$$\triangle^I r_3 = \left(a^2+b^2\right) f'' + \left(c^2+d^2\right) g''. \tag{4.3}$$

Assuming $\triangle^I r_i = \lambda_i r_i$, $i = 1,2,3$, in (4.2) and (4.3) yields $\lambda_1 = \lambda_2 = 0$ and

$$\left(a^2+b^2\right) f'' + \left(c^2+d^2\right) g'' = \lambda\left(f+g\right),\ \lambda_3 = \lambda. \tag{4.4}$$

If $\lambda = 0$ in (4.4), then we derive

$$f(u) = c_1 u^2 + c_2 u + c_3$$

and

$$g(v) = -\frac{c_1\left(a^2+b^2\right)}{\left(c^2+d^2\right)} v^2 + c_4 v + c_5,\ c_1,...,c_5 \in \mathbb{R},$$

which proves the statement (i) of the theorem.

If $\lambda \neq 0$ then (4.4) can be rewritten as

$$\left(a^2+b^2\right) f'' - \lambda f = \mu = -\left(c^2+d^2\right) g'' + \lambda g,\ \mu \in \mathbb{R}. \tag{4.5}$$

In the case $\lambda > 0$, by solving (4.5) we obtain

$$\begin{cases} f(u) = c_1 \exp\left(\sqrt{\frac{\lambda}{a^2+b^2}}u\right) + c_2 \exp\left(-\sqrt{\frac{\lambda}{a^2+b^2}}u\right) + \frac{\mu}{\lambda}, \\ g(v) = c_3 \exp\left(\sqrt{\frac{\lambda}{c^2+d^2}}v\right) + c_4 \exp\left(-\sqrt{\frac{\lambda}{c^2+d^2}}v\right) - \frac{\mu}{\lambda}, \end{cases}$$

where $c_1,...,c_4 \in \mathbb{R}$. This gives the statement (ii) of the theorem.

Otherwise, i.e., $\lambda < 0$, then we derive

$$\begin{cases} f(u) = c_1 \cos\left(\sqrt{\frac{-\lambda}{a^2+b^2}}u\right) + c_2 \sin\left(\sqrt{\frac{-\lambda}{a^2+b^2}}u\right) + \frac{\mu}{\lambda}, \\ g(v) = c_3 \cos\left(\sqrt{\frac{-\lambda}{c^2+d^2}}v\right) + c_4 \sin\left(\sqrt{\frac{-\lambda}{c^2+d^2}}v\right) - \frac{\mu}{\lambda} \end{cases}$$

for $c_1,...,c_4 \in \mathbb{R}$. This completes the proof. □

Example 4.1. Take the affine translation surface of Type 1 in $\mathbb{I}^3$ with

$$z(x,y) = \cos(x+y) + \sin(x-y),\ -\pi \leq x,y \leq \pi.$$

Then it satisfies $\triangle^I r_i = \lambda_i r_i$ for $\lambda_1 = \lambda_2 = 0$, $\lambda_3 = -2$ and can be drawn as in Fig. 2.

Next, we consider the affine translation surface of Type 1 in $\mathbb{I}^3$ that satisfies $\triangle^{II} r_i = \lambda_i r_i$, $\lambda_i \in \mathbb{R}$. Then its Laplace operator with respect to the second fundamental form II has the form

$$\begin{aligned} \triangle^{II}\phi = {} & \frac{\left(f''g''\right)^{-2}}{2(ad-bc)}\left[\left(-b\phi_x + a\phi_y\right)\left(f''\right)^2 g''' + \left(d\phi_x - c\phi_y\right) f'''\left(g''\right)^2\right] \\ & + \frac{\left(f''g''\right)^{-1}}{(ad-bc)^2}\left[\left(2ab\phi_{xy} - b^2\phi_{xx} - a^2\phi_{yy}\right) f'' + \left(2cd\phi_{xy} - d^2\phi_{xx} - c^2\phi_{yy}\right) g''\right] \end{aligned} \tag{4.6}$$

for a smooth function ϕ and $f''g'' \neq 0$. Hence we have the following result.

Theorem 4.2. *Let M be an affine translation surface of Type 1 in $\mathbb{I}^3$ that satisfies $\triangle^{II} r_i = \lambda_i r_i$. Then it is congruent to one of the following surfaces:*

(i) $(\lambda_1 \neq 0, \lambda_2 \neq 0, 0)$

$$z(x,y) = \ln\left(x^{\frac{1}{\lambda_1}} y^{\frac{1}{\lambda_2}}\right) + c_1,\ c_1 \in \mathbb{R};$$

(ii) $(\lambda \neq 0, \lambda, 0)$

$$z(x,y) = \ln\left((uv)^{\frac{1}{\lambda}}\right) + c_1,\ c_1 \in \mathbb{R},$$

where (u,v) is the affine parameter coordinates given by (3.1).

Proof. Let us assume that $\triangle^{II} r_i = \lambda_i r_i$, $\lambda_i \in \mathbb{R}$. Then, from (4.1) and (4.6), we state the following system

$$d\frac{f'''}{(f'')^2} - b\frac{g'''}{(g'')^2} = 2(ad-bc)\lambda_1 x, \tag{4.7}$$

$$-c\frac{f'''}{(f'')^2} + a\frac{g'''}{(g'')^2} = 2(ad-bc)\lambda_2 y, \tag{4.8}$$

$$\frac{f'''f'}{(f'')^2} + \frac{g'''g'}{(g'')^2} - 4 = 2\lambda_3(f+g). \tag{4.9}$$

In order to solve above system we have to distinguish two cases depending on the constants a, b, c, d for $ad - bc \neq 0$.

Case (a). Two of a, b, c, d are zero. Without loss of generality we may assume that $b = c = 0$ and $a = d = 1$. Then the equations (4.7) and (4.8) reduce to

$$\frac{f'''}{(f'')^2} = 2\lambda_1 x \tag{4.10}$$

and

$$\frac{g'''}{(g'')^2} = 2\lambda_2 y. \tag{4.11}$$

If $\lambda_1 = \lambda_2 = 0$ then we obtain a contradiction from (4.9) since f, g are non-constant functions. Thereby we need to consider the remaining cases:

Case (a.1). $\lambda_1 = 0$, i.e. $f''' = 0$. Then substituting (4.10) and (4.11) into (4.9) implies $\lambda_3 = 0$ and

$$g(y) = \frac{2}{\lambda_2}\ln y + c_1,\ c_1 \in \mathbb{R}.$$

Substituting it in (4.11) gives a contradiction.

Case (a.2). $\lambda_2 = 0$, i.e. $g''' = 0$. Hence we can similarly obtain $\lambda_3 = 0$ and

$$f(x) = \frac{2}{\lambda_1}\ln x + c_1,\ c_1 \in \mathbb{R},$$

which gives a contradiction by considering it into (4.10).

Case (a.3). $\lambda_1\lambda_2 \neq 0$. By substituting (4.10) and (4.11) into (4.9) we deduce

$$\lambda_1 x f' + \lambda_2 y g' - 2 = \lambda_3(f+g). \tag{4.12}$$

Case (a.3.1). If $\lambda_3 = 0$, then (4.12) reduces to

$$\lambda_1 x f' + \lambda_2 y g' = 2. \tag{4.13}$$

By solving (4.13) we find

$$f(x)=\frac{\xi}{\lambda_1}\ln x+c_1 \text{ and } g(v)=\frac{2-\xi}{\lambda_2}\ln y+c_2,\ c_1,c_2\in\mathbb{R},\ \xi\in\mathbb{R}^*. \tag{4.14}$$

Substituting (4.14) into (4.10) and (4.11) yields $\xi=1$. This proves the first statement of the theorem.

Case (a.3.2). If $\lambda_3\neq 0$ in (4.12) then we can rewrite it as

$$\lambda_1 x f'-\lambda_3 f-2=c_1=-\lambda_2 y g'+\lambda_3 g,\ c_1\in\mathbb{R}. \tag{4.15}$$

After solving (4.15) , we conclude

$$f(x)=-\frac{2+c_1}{\lambda_3}+c_2 x^{\frac{\lambda_3}{\lambda_1}} \tag{4.16}$$

and

$$g(y)=\frac{c_1}{\lambda_3}+c_3 y^{\frac{\lambda_3}{\lambda_2}},\ c_2,c_3\in\mathbb{R}. \tag{4.17}$$

By considering (4.16) and (4.17) into (4.10) and (4.11), respectively, we conclude $\lambda_3=0$, which implies that this case is not possible.

Case (b). At most one of a,b,c,d is zero. Suppose that $\lambda_1=0$ in (4.7). It follows from (4.7) that

$$\frac{f'''}{(f'')^2}=\frac{c_1}{d} \text{ and } \frac{g'''}{(g'')^2}=\frac{c_1}{b},\ c_1\in\mathbb{R}, \tag{4.18}$$

where we may assume that $b\neq 0\neq d$ since at most one of a,b,c,d can vanish. If $c_1=0$ then we derive a contradiction from (4.9) since $f''g''\neq 0$. Considering (4.18) into (4.8) yields $\frac{c_1}{bd}=2\lambda_2 y$, which is no possible since y is an independent variable. This implies that λ_1 is not zero and it can be similarly shown that λ_2 is not zero. Hence from (4.7) and (4.8) we can write

$$\frac{f'''}{(f'')^2}=2(\lambda_1 a x+\lambda_2 b y) \tag{4.19}$$

and

$$\frac{g'''}{(g'')^2}=2(\lambda_1 c x+\lambda_2 d y). \tag{4.20}$$

Compatibility condition in (4.19) or (4.20) gives $\lambda_1=\lambda_2$. Put $\lambda_1=\lambda_2=\lambda$. By substituting (4.19) and (4.20) into (4.9) we deduce

$$\lambda u f'+\lambda v g'-2=\lambda_3(f+g), \tag{4.21}$$

where (u,v) is the affine parameter coordinates given by (3.1).

Case (b.1). If $\lambda_3=0$, then (4.21) reduces to

$$\lambda u f'+\lambda v g'=2. \tag{4.22}$$

By solving (4.22) we find

$$f(u)=\frac{\xi}{\lambda}\ln u+c_1 \text{ and } g(v)=\frac{2-\xi}{\lambda}\ln v+c_2,\ c_1,c_2\in\mathbb{R},\ \xi\in\mathbb{R}^*. \tag{4.23}$$

Substituting (4.23) into (4.19) and (4.20) yields $\xi=1$. This proves the second statement of the theorem.

Case (b.2). If $\lambda_3\neq 0$ in (4.11) then we can rewrite it as

$$\lambda u f'-\lambda_3 f-2=c_1=-\lambda v g'+\lambda_3 g,\ c_1\in\mathbb{R}. \tag{4.24}$$

After solving (4.24) , we deduce

$$f(u) = -\frac{2+c_1}{\lambda_3} + c_2 u^{\frac{\lambda_3}{\lambda}} \tag{4.25}$$

and

$$g(v) = \frac{c_1}{\lambda_3} + c_3 v^{\frac{\lambda_3}{\lambda}},\ c_2, c_3 \in \mathbb{R}. \tag{4.26}$$

Considering (4.25) and (4.26) into (4.19) and (4.20), respectively, we find $\lambda_3 = 0$, however this is a contradiction.

□

Example 4.2. Given the affine translation surface of Type 1 in $\mathbb{I}^3$ as follows

$$z(x,y) = \ln(2x+y) + \ln(x-y), (u,v) \in [3,5] \times [1,2].$$

Then it holds $\triangle^{II} r_i = \lambda_i r_i$ for $(\lambda_1, \lambda_2, \lambda_3) = (1,1,0)$ and we plot it as in Fig. 3.



FIGURE 1. A (linear) Weingarten affine translation surface of Type 1.



FIGURE 2. An affine translation surface of Type 1 with $\triangle^I r_i = \lambda_i r_i$, $(\lambda_1, \lambda_2, \lambda_3) = (0,0,2)$.

REFERENCES

[1] M.E. Aydin, I. Mihai, *On certain surfaces in the isotropic 4-space*, Math. Commun., in press.

[2] M.E. Aydin, A. Ogrenmis, *Homothetical and translation hypersurfaces with constant curvature in the isotropic space*, In: Proceedings of the Balkan Society of Geometers, vol 23, pp.1-10, (2015).

[3] M. E. Aydin, *A generalization of translation surfaces with constant curvature in the isotropic space*, J. Geom. **107(3)** (2016), 603-615.



FIGURE 3. An affine translation surface of Type 1 with $\triangle^{II} r_i = \lambda_i r_i$, $(\lambda_1, \lambda_2, \lambda_3) = (1, 1, 0)$.

[4] Ch. Baba-Hamed, M. Bekkar, H. Zoubir, *Translation surfaces in the three-dimensional Lorentz-Minkowski space satisfying* $\triangle r_i = \lambda_i r_i$, Int. J. Math. Anal. **4(17)** (2010), 797 - 808.

[5] M. Bekkar, B. Senoussi, *Translation surfaces in the 3-dimensional space satisfying* $\triangle^{II} r_i = \mu_i r_i$, J. Geom. **103(3)** (2012), 367-374.

[6] B. Bukcu, D.W. Yoon, M.K. Karacan, *Translation surfaces in the 3-dimensional simply isotropic space* $\mathbb{I}_3^1$ *satisfying* $\triangle^{III} x_i = \lambda_i x_i$, Konuralp J. Math. **4(1)** (2016), 275-281.

[7] M. Cetin, Y. Tuncer, N. Ekmekci, *Translation surfaces in Euclidean 3-space*, Int. J. Phys. Math. Sci. **2** (2011), 49-56.

[8] B.- Y. Chen, *On submanifolds of finite type*, Soochow J. Math. **9** (1983), 65-81.

[9] B. Y. Chen, Total Mean Curvature and Submanifolds of Finite Type, World Scientific, Singapor-New Jersey-London, 1984.

[10] B. Y. Chen, S. Decu, L. Verstraelen, *Notes on isotropic geometry of production models*, Kragujevac J. Math. **7(2)** (2013), 217–220.

[11] B. Y. Chen, *Some open problems and conjectures on submanifolds of finite type: recent development*, Tamkang. J. Math. **45(1)** (2014), 87-108.

[12] F. Dillen, L. Verstraelen, G. Zafindratafa, *A generalization of the translation surfaces of Scherk*, Differential Geometry in Honor of Radu Rosca: Meeting on Pure and Applied Differential Geometry, Leuven, Belgium, 1989, KU Leuven, Departement Wiskunde (1991), pp. 107–109.

[13] M.K. Karacan, D.W. Yoon, B. Bukcu, *Translation surfaces in the three dimensional simply isotropic space* $\mathbb{I}_3^1$, Int. J. Geom. Methods Mod. Phys. **13**, 1650088 (2016) (9 pages) DOI: http://dx.doi.org/10.1142/S0219887816500882.

[14] H. Liu, Y. Yu, *Affine translation surfaces in Euclidean 3-space*, In: Proceedings of the Japan Academy, Ser. A, Mathematical Sciences, vol. **89**, pp. 111–113, Ser. A (2013).

[15] H. Liu, S.D. Jung, *Affine translation surfaces with constant mean curvature in Euclidean 3-space*, J. Geom., DOI 10.1007/s00022-016-0348-9, in press.

[16] R. Lopez , M. Moruz, *Translation and homothetical surfaces in Euclidean space with constant curvature*, J. Korean Math. Soc. **52(3)** (2015), 523-535.

[17] Z. Milin-Sipus, B. Divjak, *Mappings of ruled surfaces in simply isotropic space* I_3^1 *that preserve the generators*, Monatsh. Math. **139** (2003), 235–245.

[18] Z. Milin-Sipus, *Translation surfaces of constant curvatures in a simply isotropic space*, Period. Math. Hung. **68** (2014), 160–175.

[19] M. Moruz, M.I. Munteanu, *Minimal translation hypersurfaces in* E^4, J. Math. Anal. Appl. **439(2)** (2016), 798-812.

[20] M.I. Munteanu, O. Palmas, G. Ruiz-Hernandez, *Minimal translation hypersurfaces in Euclidean spaces*, Mediterr. J. Math. **13** (2016), 2659–2676.

[21] A.O. Ogrenmis, *Rotational surfaces in isotropic spaces satisfying Weingarten conditions*, Open Physics **14(9)** (2016), 221–225.

[22] H. Pottmann, P. Grohs, N.J. Mitra, *Laguerre minimal surfaces, isotropic geometry and linear elasticity*, Adv. Comput. Math. **31** (2009), 391–419.

[23] H. Sachs, Isotrope Geometrie des Raumes, Vieweg Verlag, Braunschweig, 1990.

[24] K. Seo, *Translation Hypersurfaces with constant curvature in space forms*, Osaka J. Math. **50** (2013), 631-641.
[25] K. Strubecker, *ber die isotropen Gegenstcke der Minimalche von Scherk*, J. Reine Angew. Math. **293** (1977), 2251.
[26] L. Verstraelen, J. Walrave, S. Yaprak, *The minimal translation surfaces in Euclidean space*, Soochow J. Math. **20** (1994), 77–82.

* Department of Mathematics, Faculty of Science, Firat University, Elazig, 23200, Turkey

Department of Mathematics, Faculty of Science and Art, Namik Kemal University, Tekirdag, 59000, Turkey
E-mail address: meaydin@firat.edu.tr, mergut@nku.edu.tr